

Transcriptional activation by AP-2 α is modulated by the oncogene DEK

Mónica Campillos, Miguel Angel García, Fernando Valdivieso and Jesús Vázquez*

Centro de Biología Molecular 'Severo Ochoa', CSIC-Universidad Autónoma de Madrid, 28049 Cantoblanco, Madrid, Spain

Received September 11, 2002; Revised November 15, 2002; Accepted January 12, 2003

ABSTRACT

Cell differentiation and development are highly regulated processes at the transcriptional level. One of the main transcription factors that regulate these processes is AP-2 α , a cell-type specific protein required for vertebrate development and embryogenesis. AP-2 α also regulates apoptosis and cell-cycle specific events by interacting with the oncogene c-Myc. In searching for novel AP-2 α -interacting factors, using an affinity chromatography approach, we have observed that oncoprotein DEK interacts with AP-2 α *in vitro*. The existence of an interaction between AP-2 α and DEK in cellular cultures was demonstrated by expression of a tagged AP-2 α form followed by immunodetection. By transient co-expression experiments using a reporter for *APOE* promoter activity we have found that DEK stimulates the transactivation activity of AP-2 α over *APOE* promoter. Finally, electrophoretic mobility shift assays suggested that DEK enhances the DNA-binding activity of AP-2 α . Our data suggest a novel cellular function of DEK as a transcriptional co-activator.

INTRODUCTION

The family of AP-2 transcription factors is composed of four members: AP-2 α (1,2), AP-2 β (3), AP-2 γ (4) and AP-2 δ (5). Spatially and temporally restricted expression patterns of these proteins (6–8) provide essential clues for embryonic development (9,10), regulation of programmed cell death (11), and cell growth and differentiation (12).

Recently, the relevance of protein–protein interaction on AP-2 α activity has been reported. AP-2 α modulates the function of Myc as a proliferative agent and also as apoptosis inducer (13). Other known AP-2 α -interacting proteins are the retinoblastoma Rb protein (14,15), the Yin Yang 1 factor (YY1) (16), and the transcription factors YB-1 (17) and Sp-1 (18).

We have recently described the presence of allelic polymorphisms in the transcriptional regulatory region of the *APOE* gene, which produce variations in promoter activity

(19). Two of these variants are associated with an increase in risk for developing late-onset Alzheimer's disease (20,21). We also found that the activity of the proximal *APOE* promoter is upregulated by cAMP and retinoic acid, in astrocytic but not hepatic cells (22). The cAMP effect is mediated, in part, by interaction of factor AP-2 α with two sites located in the *APOE* proximal region (22). The stimulatory effect of cAMP on *APOE* promoter in HepG2 cells involves AP-2 α phosphorylation at Ser239 by protein kinase A (23). In this work, we investigate the existence of potential AP-2 α -interacting factors which may modulate AP-2 α activity. We have identified the oncoprotein DEK as an AP-2 α -binding factor; DEK was found to enhance the effect of AP-2 α on *APOE* promoter, probably enhancing the DNA binding of AP-2 α .

MATERIALS AND METHODS

Recombinant protein expression

For the construction of the pcDNA3hisAP-2 vector, which expresses an N-terminal truncation of human AP-2 α (deletion of the first 122 amino acids) with a poly-histidine extension at its N-terminal end (hisAP-2), a fragment was amplified by PCR using pTrcHisBAP-2 vector (23) as template, followed by cloning of the fragment into the pcDNA3 vector using the *Xho*I and *Hind*III cloning sites.

For purification of HisAP-2 from U-87 derived cell lines, identical amounts of nuclear protein from 25×10^6 cells (24) were diluted 50-fold with buffer D (0.2 M KCl, 20 mM Tris–HCl, pH 7.6, 10% glycerol, 2 mM MgCl₂ and protease inhibitors) and loaded onto a 200 μ l Ni²⁺ column (Invitrogen Corporation) previously equilibrated in the same buffer. The column was washed with 20 vol of buffer D and eluted in 200 μ l fractions with buffer D containing 500 mM imidazole.

Human recombinant AP-2 α (AP-2r) was prepared as described previously (23). GST-DEK was an N-terminally truncated human DEK form containing 80 amino acids fused to GST kindly provided by Gerald Grosveld.

AP-2r affinity chromatography

Nuclear extracts from rat liver were prepared as described previously (25). The extracts were dialyzed against buffer B (20 mM Tris–HCl pH 7.6, 10% glycerol, 2 mM MgCl₂, 0.2 mM EDTA and 1 mM DTT) containing 0.2 M KCl and then passed through 30 ml of a heparin–Sepharose column

*To whom correspondence should be addressed. Tel: +34 913 978364; Fax: +34 913 978087; Email: jvazquez@cbm.uam.es

The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors




Figure 1. Purification of DEK by AP-2 α affinity chromatography. A rat nuclear extract (E) was fractionated and enriched using an AP-2 α affinity column (see Materials and Methods). The eluted fractions were pooled and subjected to SDS-PAGE, followed by silver staining (1). The band indicated by an arrow was identified as the rat homologue of human DEK by mass spectrometry.

(Amersham Pharmacia Biotech). The 0.5 M KCl buffer B eluted fractions were pooled, dialyzed against 0.1 M KCl buffer C (25 mM Tris-HCl pH 7.6, 12.5 mM MgCl₂, 20% glycerol, 1 mM DTT and 0.1% NP-40) and passed through an AP-2r-agarose column. This column was eluted with 1 M KCl buffer C. The AP-2r affinity column was constructed by coupling 250 mg of agarose (CNBr-activated Sepharose 4B) to 8 mg AP-2r (23) following the manufacturer's instructions (Amersham Pharmacia Biotech).

Cell lines, transfections, luciferase and β -galactosidase assays and electrophoretic mobility shift assays (EMSAs)

The cell lines U-87 MG (ATCC number: HTB-14) and HepG2 (ATCC number: HB-8065) were grown and transfected as described (23). For the selection of stable U-87 cell line expressing hisAP-2 protein, 10⁶ cells were transfected with 20 μ g of the pcDNA3hisAP2 vector and grown in selective medium (geneticin 0.4 mg/ml). After 2 days cells were plated in a limiting dilution in 24-well plates and AP-2 α expression in single colonies was analysed by western blot using the anti-humanAP-2 antibody C-18 (Santa Cruz, CA). Luciferase and β -galactosidase activities were measured as described previously (23). EMSAs were carried out as reported (23). The CXX oligonucleotide sequence used in this work was the following: 5'-CTGTGCCTGGGGCAGGGGGAGAACA-3'.

Mass spectrometry

The identity of the purified band was determined following the same procedure described previously (23).

RESULTS

Identification of DEK as an AP-2 α interacting factor using AP-2 α -affinity chromatography

To identify potential AP-2 α -interacting proteins, we carried out a two-step AP-2 α -affinity chromatography purification using nuclear extracts from rat liver. A 50 kDa protein band was clearly enriched in the eluted fraction (Fig. 1). This band was excised from a silver stained SDS-polyacrylamide gel, subjected to in-gel tryptic cleavage and analysed by nanospray-ion trap MS/MS mass spectrometry. From the




Figure 2. Interaction of hisAP-2 with human DEK in U-87, D7 and D10 astrocytoma cell lines. (A) Analysis of the expression of hisAP-2 in the stably transfected clones. Nuclear extracts (20 μ g) from U-87, D7 and D10 were analysed by western blotting using C-18 antibody. As a positive control we used 1 μ g of AP-2r. The arrow indicates the position of hisAP-2 band. (B) Purification of hisAP-2 from the stably transfected cell clones. Purified fractions (30 μ l) from U-87, D7 and D10 were analysed by western blotting using the C-18 antibody. The arrow indicates position of hisAP-2. (C) DNA-binding activity of the purified fractions. EMSAs were carried out using 5 and 15 μ l of the purified fractions from U-87, D7 and D10 and CXX probe. As a positive control 0.5 and 1 μ g of AP-2r were used. The arrow indicates the position of the hisAP-2-DNA complex. (D) Co-purification of DEK with AP-2 α . Ten micrograms of the original nuclear extracts and 40 μ g of purified fractions from U-87, D7 and D10 were analysed by western blotting using a polyclonal antibody that recognizes the human DEK protein (33). The arrow indicates the position of the DEK band. In (A), (B) and (D), the molecular weight markers are indicated on the left.

total mixture of tryptic peptides, three of them (EPFTIAQGK, LLTNRPGTVSSLK and NVGQFSGFPFEK) were identified by a database search. All sequenced peptides showed a 100% identity to the human DEK protein, a 43 kDa protein initially discovered as a fusion partner of the CAN nucleoporin in a specific subtype of acute myeloid leukaemia (26). The identity between the sequenced peptides and the sequence of human DEK indicates that the purified protein is the DEK rat homologue, which has still not been cloned. We then investigated whether a physiologically relevant interaction between AP-2 α and DEK occurs in human cells.

The oncoprotein DEK interacts with AP-2 α in human cells

An N-terminal histidine-tagged AP-2 α protein (hisAP-2) was stably expressed in the astrocytoma cell line U-87. Two highly expressing hisAP-2 U-87 clones, D7 and D10, were selected (Fig. 2A, lanes 2 and 3). From these two cell clones, we were only able to purify hisAP-2 from protein extracts from the D10 clone by using Ni²⁺ columns (Fig. 2B, lane 3), suggesting that the form of hisAP-2 expressed by D7 contained a




Figure 3. Effect of DEK on the transcriptional activity of AP-2 α in HepG2 cells. 300 ng of the *APOE* promoter luciferase vector pXP2-227 were co-transfected with 300 ng of pcDNA3 (grey bars) or with an AP-2 α expression vector (black bars) and increasing quantities of 0, 200, 400, 600 or 800 ng of the DEK expression vector combined with decreasing quantities of pcDNA3 of 800, 600, 400, 200 and 0 ng. Luciferase activity was corrected for the transfection efficiency taking into account the activity of 300 ng of β -galactosidase expression vector. Transfection assays were performed in triplicate and results are representative of at least three independent experiments; data are expressed as mean \pm SEM.

non-functional protein due to some mutation or deletion in the Ni²⁺-binding domain. Consistently, the fractions purified from protein extracts from D7 did not present any noticeable DNA binding activity, whereas this activity was detected in the fractions eluted from Ni²⁺ columns loaded with clone D10 extracts (Fig. 2C, compare lanes 6 and 7 with 8 and 9). We therefore used the D7 clone as an internal control in subsequent experiments. An unidentified 110 kDa band co-purified with AP-2 α and was recognised by the anti-AP-2 antibody (Fig. 2B, lane 3). A band of similar molecular weight has been described by other authors using anti-AP-2 α antibodies (27,28).

We next investigated whether DEK co-purified with hisAP-2. The nuclear expression levels of DEK in the two clones, D7 and D10, were similar to those of the original U-87 cell line (Fig. 2D, lanes 1–3). DEK was clearly detected in the AP-2 α -enriched fractions when nuclear extracts from clone D10 were subjected to Ni²⁺-affinity chromatography (Fig. 2D, lane 6). In clear contrast, DEK was absent in the eluates from either U-87 or clone D7 (Fig. 2D, lanes 4 and 5). These data strongly suggest that DEK interacts with AP-2 α in cell cultures.

DEK increases transcriptional activity of AP-2 α in HepG2 cells

To investigate the physiological relevance of the DEK-AP-2 α interaction, we tested whether DEK could modulate AP-2 α transcriptional activity. We co-transfected increasing amounts of a DEK expression vector with a constant amount of an AP-2 α expression vector and an *APOE* promoter-luciferase reporter vector [from –227 to +1 *APOE* promoter sequence (22)] in the hepatoma cell line HepG2. According to previously reported results (22,23,29), AP-2 α transactivated *APOE* promoter in these cells (Fig. 3). The addition of increasing amounts of DEK enhanced AP-2 α activity in a dose-dependent manner (Fig. 3, black bars). AP-2 α was directly involved in this effect, since overexpression of DEK




Figure 4. Effect of DEK on binding of AP-2r to DNA *in vitro*. CXX probe was incubated with 10 (lanes 2–5), 30 (lanes 6–9) or 100 ng (lanes 10–13) of AP-2r in the presence of either 20 (lanes 3, 7 and 11), 100 (lanes 4, 8 and 12) or 500 ng (lanes 5, 9 and 13) of GST-DEK. As negative controls, CXX probe was incubated with 100 ng of AP-2r in the presence of either 20 (lane 16), 100 (lane 17) or 500 ng (lane 18) of GST, and 500 ng of GST-DEK (lane 14) and GST (lane 19) were incubated with the probe in the absence of AP-2r.

alone did not have any effect on *APOE* promoter activity (Fig. 3, grey bars). Taken together, these results suggested that DEK plays a role as co-activator of the transcriptional activity of AP-2 α .

DEK enhances binding of AP-2 α to DNA

We then examined whether a recombinant preparation of GST-DEK affected the formation of AP-2 α -DNA complexes by EMSAs. As shown in Figure 4, while *in vitro* DNA binding activity of small amounts of AP-2r (30 ng) was undetectable, formation of AP-2r-DNA complexes could be observed in the presence of GST-DEK. This effect of GST-DEK appeared to be specific since GST did not enhance the DNA-binding activity of AP-2 α (Fig. 4). The stimulatory effect of GST-DEK was dose dependent (Fig. 4), and we were unable to observe the formation of DNA complexes with GST-DEK alone (Fig. 4). In addition, the presence of GST-DEK did not produce a significant increase in the mobility of the AP-2 α -DNA complex. In our assay conditions we have been unable to detect a supershift band with an anti-human DEK antibody (30) (data not shown) suggesting that DEK is not present in the complex; it is also possible that DEK is not recognised by the antibody in the presence of bound AP-2 α due to some kind of steric hindrance. Taken together, our results suggest the existence of a physiologically relevant interaction between the oncoprotein DEK and the transcription factor AP-2 α ; DEK enhances AP-2 α transcriptional activity by a mechanism that appears to involve an enhancement in AP-2 α -DNA binding.

DISCUSSION

To the best of our knowledge, this is the first report that suggests a role for the oncogene DEK as co-activator of a transcription factor. DEK contains a SAP (after SAF-A/B, Acinus and PIAS) DNA-binding motif which is involved in chromatin remodelling and transcription (31). Coincidentally,

several SAP family members, such as Miz1 (32), alter the DNA binding activity of transcription factors, modifying their transcriptional activity. Since DEK is known to bind condensed chromosomes (33) and chromatin (30,34), altering the nucleosomal DNA topology and reducing the chromatin replication efficiency (35), it is possible that DEK produces a change in chromatin conformation that increases the DNA binding of AP-2 α . The AP-2 α binding site on *APOE* promoter (TGGGGCAGGG) is very similar to the known DEK binding site in the HIV-2 promoter, which is a *pac* sequence (TTGGTCAGGG) (30). Although DEK does not appear to bind to the CXX probe *in vitro*, it is possible that, upon a possible *in vivo* binding to DNA, DEK may act as a recruitment factor for AP-2 α . In *Saccharomyces cerevisiae*, the expression of Pho5 is controlled by a similar 'cooperativity' mechanism. In this case, the Pho4 transcriptional activator is cooperatively recruited to the active sites by the homeoprotein Pho2, a protein that binds very weakly to DNA in the absence of Pho4 (36).

However, DEK appears to enhance the interaction of AP-2 α with DNA without affecting the rate of migration of this complex, and no evidence was found suggesting a direct interaction of DEK with DNA. Thus, DEK behaves similarly to Miz1, another SAP protein, which enhances DNA binding of the transcription factor Mx2 without forming ternary complexes (32). Our results suggest the existence of a transient interaction in which DEK affects the conformation of AP-2 α , enhancing its DNA binding. A similar 'hit and run' mechanism has been demonstrated for the effect of Phox 1 over DNA binding of the serum response factor (SRF); Phox 1 was proposed to lower an activation energy barrier, enhancing the rate of stable SRF-DNA complex formation (37).

The interaction of AP-2 α with DEK was demonstrated using an N-terminally truncated AP-2 α fragment. This suggests that the interaction domain of AP-2 α is located in the C-terminal moiety, which includes the DNA binding and dimerization domains. Both domains are highly conserved between all members of the AP-2 family (5,38) suggesting that DEK may be interacting with other AP-2 family members.

It has recently been suggested that cardiac, neural crest and neural tube defects in mice lacking Cited2 are produced because this protein plays a role as AP-2 α co-activator (39). Also, over-expression of some co-activators, such as PC4 and PARP, partially rescue the malignant phenotype of PA-1 cells transformed by AP-2 α via a mechanism that involve an effect of these co-activators over other transcription factors (27,40). Since DEK can partially revert the transformation-prone phenotype of cells from patients with ataxia-telangiectasia (41), it is reasonable to think that DEK may rescue this phenotype due to the action of DEK-dependent transcription factors, such as AP-2 α . In this sense, we can speculate that the oncogenic activity of the chimera DEK-CAN (26) may be due to the inhibition of the interaction with AP-2 α and other transcription factors, provoking, as a result, cell transformation.

It has been demonstrated that DEK is an oncogene associated with human hepatocellular carcinogenesis (42). The DEK oncogenic properties and the co-expression of DEK and AP-2 α in several tissues such as spleen, thymus, kidney and brain (6,26) suggest that DEK could be acting in a coordinate way with AP-2 α in the cell cycle regulation.

ACKNOWLEDGEMENTS

We are very grateful to Professor Gerald Grosveld for the generous gifts of GST-DEK recombinant protein and polyclonal antibody against human DEK, and for helpful comments on the manuscript. We also thank the anonymous reviewers whose suggestions helped to clarify this work. This work was supported by CICYT SAF 2000-0178 and 08.5/0065.1/2001 grants from Spanish Ministerio de Ciencia y Tecnología and from Comunidad Autónoma de Madrid, and by an institutional grant by Fundación Ramón Areces to CBMSO.

REFERENCES

- Williams, T., Admon, A., Luscher, B. and Tjian, R. (1988) Cloning and expression of AP-2, a cell-type-specific transcription factor that activates inducible enhancer elements. *Genes Dev.*, **2**, 1557-1569.
- Mitchell, P.J., Wang, C. and Tjian, R. (1987) Positive and negative regulation of transcription *in vitro*: enhancer-binding protein AP-2 is inhibited by SV40 T antigen. *Cell*, **50**, 847-861.
- Moser, M., Imhof, A., Pscherer, A., Bauer, R., Amselgruber, W., Sinowatz, F., Hofstadter, F., Schule, R. and Buettner, R. (1995) Cloning and characterization of a second AP-2 transcription factor: AP-2 β . *Development*, **121**, 2779-2788.
- Oulad-Abdelghani, M., Bouillet, P., Chazaud, C., Dolle, P. and Chambon, P. (1996) AP-2.2: a novel AP-2-related transcription factor induced by retinoic acid during differentiation of P19 embryonal carcinoma cells. *Exp. Cell Res.*, **225**, 338-347.
- Zhao, F., Satoda, M., Licht, J.D., Hayashizaki, Y. and Gelb, B.D. (2001) Cloning and characterization of a novel mouse AP-2 transcription factor, AP-2 δ , with unique DNA binding and transactivation properties. *J. Biol. Chem.*, **276**, 40755-40760.
- Mitchell, P.J., Timmons, P.M., Hebert, J.M., Rigby, P.W. and Tjian, R. (1991) Transcription factor AP-2 is expressed in neural crest cell lineages during mouse embryogenesis. *Genes Dev.*, **5**, 105-119.
- Chazaud, C., Oulad-Abdelghani, M., Bouillet, P., Decimo, D., Chambon, P. and Dolle, P. (1996) AP-2.2, a novel gene related to AP-2, is expressed in the forebrain, limbs and face during mouse embryogenesis. *Mech. Dev.*, **54**, 83-94.
- Moser, M., Ruschhoff, J. and Buettner, R. (1997) Comparative analysis of AP-2 α and AP-2 β gene expression during murine embryogenesis. *Dev. Dyn.*, **208**, 115-124.
- Schorle, H., Meier, P., Buchert, M., Jaenisch, R. and Mitchell, P.J. (1996) Transcription factor AP-2 essential for cranial closure and craniofacial development. *Nature*, **381**, 235-238.
- Zhang, J., Hagopian-Donaldson, S., Serbedzija, G., Elsemore, J., Plehn-Dujowich, D., McMahon, A.P., Flavell, R.A. and Williams, T. (1996) Neural tube, skeletal and body wall defects in mice lacking transcription factor AP-2. *Nature*, **381**, 238-241.
- Moser, M., Pscherer, A., Roth, C., Becker, J., Mucher, G., Zerres, K., Dixkens, C., Weis, J., Guay-Woodford, L., Buettner, R. and Fassler, R. (1997) Enhanced apoptotic cell death of renal epithelial cells in mice lacking transcription factor AP-2 β . *Genes Dev.*, **11**, 1938-1948.
- Byrne, C., Tainsky, M. and Fuchs, E. (1994) Programming gene expression in developing epidermis. *Development*, **120**, 2369-2383.
- Eilers, M., Schirm, S. and Bishop, J.M. (1991) The MYC protein activates transcription of the α -prothymosin gene. *EMBO J.*, **10**, 133-141.
- Batsche, E., Muchardt, C., Behrens, J., Hurst, H.C. and Cremisi, C. (1998) RB and c-Myc activate expression of the E-cadherin gene in epithelial cells through interaction with transcription factor AP-2. *Mol. Cell. Biol.*, **18**, 3647-3658.
- Wu, F. and Lee, A.S. (1998) Identification of AP-2 as an interactive target of Rb and a regulator of the G1/S control element of the hamster histone H3.2 promoter. *Nucleic Acids Res.*, **26**, 4837-4845.
- Wu, F. and Lee, A.S. (2001) YY1 as a regulator of replication-dependent hamster histone H3.2 promoter and an interactive partner of AP-2. *J. Biol. Chem.*, **276**, 28-34.
- Mertens, P.R., Alfonso-Jaume, M.A., Steinmann, K. and Lovett, D.H. (1998) A synergistic interaction of transcription factors AP2 and YB-1

- regulates gelatinase A enhancer-dependent transcription. *J. Biol. Chem.*, **273**, 32957–32965.
18. Pena, P., Reutens, A.T., Albanese, C., D'amico, M., Watanabe, G., Donner, A., Shu, I.W., Williams, T. and Pestell, R.G. (1999) Activator protein-2 mediates transcriptional activation of the CYP11A1 gene by interaction with Sp1 rather than binding to DNA. *Mol. Endocrinol.*, **13**, 1402–1416.
 19. Artiga, M.J., Bullido, M.J., Sastre, I., Recuero, M., García, M.A., Aldudo, J., Vázquez, J. and Valdivieso, F. (1998) Allelic polymorphisms in the transcriptional regulatory region of apolipoprotein E gene. *FEBS Lett.*, **421**, 105–108.
 20. Artiga, M.J., Bullido, M.J., Frank, A., Sastre, I., Recuero, M., García, M.A., Lendon, C.L., Han, S.W., Morris, J.C., Vázquez, J., Goate, A. and Valdivieso, F. (1998) Risk for Alzheimer's disease correlates with transcriptional activity of the APOE gene. *Hum. Mol. Genet.*, **7**, 1887–1892.
 21. Bullido, M.J., Artiga, M.J., Recuero, M., Sastre, I., García, M.A., Aldudo, J., Lendon, C., Han, S.W., Morris, J.C., Frank, A., Vázquez, J., Goate, A. and Valdivieso, F. (1998) A polymorphism in the regulatory region of APOE associated with risk for Alzheimer's dementia. *Nature Genet.*, **18**, 69–71.
 22. García, M.A., Vázquez, J., Giménez, C., Valdivieso, F. and Zafra, F. (1996) Transcription factor AP-2 regulates human apolipoprotein E gene expression in astrocytoma cells. *J. Neurosci.*, **16**, 7550–7556.
 23. García, M.A., Campillos, M., Marina, A., Valdivieso, F. and Vázquez, J. (1999) Transcription factor AP-2 activity is modulated by protein kinase A-mediated phosphorylation. *FEBS Lett.*, **444**, 27–31.
 24. Andrews, N.C. and Faller, D.V. (1991) A rapid micropreparation technique for extraction of DNA-binding proteins from limiting numbers of mammalian cells. *Nucleic Acids Res.*, **19**, 2499.
 25. Zabala, M.T. and García-Ruiz, J.P. (1989) Regulation of expression of the messenger ribonucleic acid encoding the cytosolic form of phosphoenolpyruvate carboxykinase in liver and small intestine of lactating rats. *Endocrinology*, **125**, 2587–2593.
 26. Von Lindern, M., Fornerod, M., Van Baal, S., Jaegle, M., De Wit, T., Buijs, A. and Grosveld, G. (1992) The translocation (6;9), associated with a specific subtype of acute myeloid leukemia, results in the fusion of two genes, *dek* and *can*, and the expression of a chimeric, leukemia-specific *dek-can* mRNA. *Mol. Cell. Biol.*, **12**, 1687–1697.
 27. Kannan, P. and Tainsky, M.A. (1999) Coactivator PC4 mediates AP-2 transcriptional activity and suppresses ras-induced transformation dependent on AP-2 transcriptional interference. *Mol. Cell. Biol.*, **19**, 899–908.
 28. Boshier, J.M., Williams, T. and Hurst, H.C. (1995) The developmentally regulated transcription factor AP-2 is involved in c-erbB-2 overexpression in human mammary carcinoma. *Proc. Natl Acad. Sci. USA*, **92**, 744–747.
 29. García, M.A., Campillos, M., Ogueta, S., Valdivieso, F. and Vázquez, J. (2000) Identification of amino acid residues of transcription factor AP-2 involved in DNA binding. *J. Mol. Biol.*, **301**, 807–816.
 30. Fu, G.K., Grosveld, G. and Markovitz, D.M. (1997) DEK, an autoantigen involved in a chromosomal translocation in acute myelogenous leukemia, binds to the HIV-2 enhancer. *Proc. Natl Acad. Sci. USA*, **94**, 1811–1815.
 31. Aravind, L. and Koonin, E.V. (2000) SAP—a putative DNA-binding motif involved in chromosomal organization. *Trends Biochem. Sci.*, **25**, 112–114.
 32. Wu, L., Wu, H., Ma, L., Sangiorgi, F., Wu, N., Bell, J.R., Lyons, G.E. and Maxson, R. (1997) Miz1, a novel zinc finger transcription factor that interacts with Mx2 and enhances its affinity for DNA. *Mech. Dev.*, **65**, 3–17.
 33. Fornerod, M., Boer, J., Van Baal, S., Jaegle, M., Von Lindern, M., Murti, K.G., Davis, D., Bonten, J., Buijs, A. and Grosveld, G. (1995) Relocation of the carboxyterminal part of CAN from the nuclear envelope to the nucleus as a result of leukemia-specific chromosome rearrangements. *Oncogene*, **10**, 1739–1748.
 34. Faulkner, N.E., Hilfinger, J.M. and Markovitz, D.M. (2001) Protein phosphatase 2A activates the HIV-2 promoter through enhancer elements that include the *pets* site. *J. Biol. Chem.*, **276**, 25804–25812.
 35. Alexiadis, V., Waldmann, T., Andersen, J., Mann, M., Knippers, R. and Gruss, C. (2000) The protein encoded by the proto-oncogene DEK changes the topology of chromatin and reduces the efficiency of DNA replication in a chromatin-specific manner. *Genes Dev.*, **14**, 1308–1312.
 36. Barbaric, S., Munsterkotter, M., Goding, C. and Horz, W. (1998) Cooperative Pho2-Pho4 interactions at the PHO5 promoter are critical for binding of Pho4 to UASp1 and for efficient transactivation by Pho4 at UASp2. *Mol. Cell. Biol.*, **18**, 2629–2639.
 37. Grueneberg, D.A., Natesan, S., Alexandre, C. and Gilman, M.Z. (1992) Human and *Drosophila* homeodomain proteins that enhance the DNA-binding activity of serum response factor. *Science*, **257**, 1089–1095.
 38. Wankhade, S., Yu, Y., Weinberg, J., Tainsky, M.A. and Kannan, P. (2000) Characterization of the activation domains of AP-2 family transcription factors. *J. Biol. Chem.*, **275**, 29701–29708.
 39. Bamforth, S.D., Braganca, J., Eloranta, J.J., Murdoch, J.N., Marques, F.I., Kranc, K.R., Farza, H., Henderson, D.J., Hurst, H.C. and Bhattacharya, S. (2001) Cardiac malformations, adrenal agenesis, neural crest defects and exencephaly in mice lacking Cited2, a new Tfp2 co-activator. *Nature Genet.*, **29**, 469–474.
 40. Kannan, P., Yu, Y., Wankhade, S. and Tainsky, M.A. (1999) PolyADP-ribose polymerase is a coactivator for AP-2-mediated transcriptional activation. *Nucleic Acids Res.*, **27**, 866–874.
 41. Meyn, M.S., Lu-Kuo, J.M. and Herzing, L.B. (1993) Expression cloning of multiple human cDNAs that complement the phenotypic defects of ataxia-telangiectasia group D fibroblasts. *Am. J. Hum. Genet.*, **53**, 1206–1216.
 42. Kondoh, N., Wakatsuki, T., Ryo, A., Hada, A., Aihara, T., Horiuchi, S., Goseki, N., Matsubara, O., Takenaka, K., Shichita, M., Tanaka, K., Shuda, M. and Yamamoto, M. (1999) Identification and characterization of genes associated with human hepatocellular carcinogenesis. *Cancer Res.*, **59**, 4990–4996.